



# Construction of Actuarial Models

Fourth Edition

by Mike Gauger and Michael Hosking

Published by BPP Professional Education

## Solutions to practice questions – Chapter 13

### Solution 13.1

Using the upper 5% point of the normal distribution, we have:

$$Q_{0.95} = 500 + 1.6449\sqrt{100} = 516.449$$

### Solution 13.2

The normal distribution will have a thinner right hand tail than the corresponding lognormal distribution.

For large claim amounts, the lognormal curve will lie above the corresponding normal curve.

So if we want to contain an area to the right equal to 5%, we will need to move our risk measure to the right, *ie* it should be a little bigger. Let's now check this out.

First we need the parameters of the lognormal distribution:

$$e^{\mu + \frac{1}{2}\sigma^2} = 500 \qquad e^{2\mu + \sigma^2} (e^{\sigma^2} - 1) = 100$$

Solving these simultaneous equations, we find that  $\mu = 6.214408$  and  $\sigma^2 = 0.00039992$ .

We now need the 95<sup>th</sup> percentile of this distribution:

$$\Pr(\log N(\mu, \sigma^2) < Q) = \Pr(N(\mu, \sigma^2) < \log Q) = \Pr\left(N(0, 1) < \frac{\log Q - \mu}{\sigma}\right) = 0.95$$

This gives us the equation:

$$\frac{\log Q - 6.214408}{\sqrt{0.00039992}} = 1.6449 \quad \Rightarrow \quad Q = 516.62$$

As we anticipated, we have a slightly greater risk measure.

**Solution 13.3**

We need to solve the equation:

$$F(x) = 1 - \left( \frac{\theta}{\theta + x} \right)^\alpha = 0.95$$

Substituting in  $\theta = 50$  and  $\alpha = 2.5$  and rearranging, we find that:

$$x_{0.95} = \frac{50}{0.05^{0.4}} - 50 = 115.723$$


---

**Solution 13.4**

Using the fact that  $X - x_\alpha | X > x_\alpha$  is also Pareto, but with parameters  $\theta' = 50 + 115.723$  and  $\alpha = 2.5$ , we see that the value of the 95% CTE risk measure in this case is:

$$x_\alpha + \frac{\theta'}{\alpha - 1} = 115.723 + \frac{50 + 115.723}{1.5} = 226.205$$


---

**Solution 13.5**

First we need the VaR risk measure. Using the distribution function of the exponential distribution:

$$F(x) = 0.98 = 1 - e^{-x/500} \Rightarrow x_\alpha = -500 \log 0.02 = 1,956.01$$

The exponential distribution also has a useful conditional distribution result. If  $X$  has an exponential distribution with mean  $\theta$ , then  $X - x_\alpha | X > x_\alpha$  also has an exponential distribution with the same mean (this is the lack of memory property of the exponential distribution). So the CTE risk measure is:

$$1,956.01 + 500 = 2,456.01$$


---

**Solution 13.6**

First, we see that the CDF of the loss distribution has a discontinuity at  $x = 50$ . The distribution jumps up from  $1 - 0.082085 = 0.917915$  to 1 at that point.

So  $Q_{0.95}$  and  $Q_{0.99}$  are both equal to 50.

But the TVaR risk measures are the values of the conditional expectation, given that the loss is greater than or equal to 50. But in this case the loss must be exactly 50, since it can never exceed 50. So both TVaR risk measures are equal to 50.

**Solution 13.7**

The distribution function for the Weibull distribution is:

$$F(x) = 1 - e^{-(x/\theta)^\tau}$$

We need to solve the equation:

$$1 - e^{-(x/500)^\tau} = 0.95 \quad \Rightarrow \quad x = \sqrt{-500^2 \log 0.05} = 865.409$$


---

**Solution 13.8**

The probability of loss is:

$$S(1,000) = e^{-(1,000/\theta)^\tau} = e^{-(2)^2} = e^{-4} = 0.01832$$


---

**Solution 13.9**

First we find the probability that  $X = 0$ :

$$\begin{aligned} \Pr(X = 0) &= \Pr(S \geq 1,000) = \Pr(1,000e^{0.04+0.2Z} \geq 1,000) = \Pr(0.04 + 0.2Z \geq 0) \\ &= \Pr(N(0.04, 0.2^2) \geq 0) = \Pr\left(N(0, 1) \geq -\frac{0.04}{0.2}\right) = 1 - \Phi(-0.2) = 0.5793 \end{aligned}$$

Since  $0.9 > 0.5793$ , the 90<sup>th</sup> percentile of the distribution of  $X$  lies in the continuous part of the distribution.

We want:

$$\begin{aligned} 0.90 &= \Pr(X \leq x_{0.90}) = \Pr(1,000 - S \leq x_{0.90}) = \Pr(1,000 - 1,000e^{0.04+0.2Z} \leq x_{0.90}) \\ &= \Pr\left(e^{0.04+0.2Z} \geq \frac{1,000 - x_{0.90}}{1,000}\right) \end{aligned}$$

But  $e^{0.04+0.2Z}$  has a lognormal distribution with  $\mu = 0.04$  and  $\sigma = 0.2$ . So we want the 10<sup>th</sup> percentile of this lognormal distribution.

The 10<sup>th</sup> percentile of  $N(0, 1)$  is  $-1.282$ . The 10<sup>th</sup> percentile of  $N(0.04, 0.2^2)$  is  $0.04 - (1.282 \times 0.2) = -0.2164$ . The 10<sup>th</sup> percentile of the corresponding lognormal distribution is  $e^{-0.2164} = 0.80541$ .

So we now have the equation:

$$\frac{1,000 - x_{0.90}}{1,000} = 0.80541$$

which gives us  $x_{0.90} = 194.59$ .

---

### Solution 13.10

First we find the relevant VaR risk measures. These are  $Q_{0.9} = 10$ ,  $Q_{0.95} = 100$  and  $Q_{0.99} = 500$ .

We now look at the outcomes for the largest 10% of probability. Of the largest 10% of probability, 0.02 of it is at 10, 0.06 is at 100, 0.016 is at 500 and 0.004 is at 1,000. Factoring these up by dividing by 0.1, we obtain the corresponding conditional distribution:

$$\Pr(X = 10) = 0.2 \quad \Pr(X = 100) = 0.6 \quad \Pr(X = 500) = 0.16 \quad \Pr(X = 1,000) = 0.04$$

The mean of this conditional distribution is :

$$10 \times 0.2 + 100 \times 0.6 + 500 \times 0.16 + 1,000 \times 0.04 = 182.$$

This is the 90% TVaR risk measure.

Alternatively we can use the formula

$$\text{TVaR}_\alpha(X) = \frac{(\beta - \alpha)x_\alpha + (1 - \beta)E[X | X > x_\alpha]}{1 - \alpha}$$

with  $\beta = 0.92$ ,  $\alpha = 0.9$ , and:

$$E[X | X > 10] = 100 \times \frac{.06}{.08} + 500 \times \frac{.016}{.08} + 1,000 \times \frac{.004}{.08} = 225$$

So now we get:

$$\text{TVaR}_{0.90} = \frac{(0.92 - 0.9)10 + (1 - 0.92)225}{0.1} = 182$$

This gives the same answer as the more intuitive approach.

You should be able to obtain (using either method) the corresponding results:

$$\text{TVaR}_{0.95} = 300 \quad \text{TVaR}_{0.99} = 700$$

**Solution 13.11**

First we find the corresponding quantile risk measures. By accumulating the probability, we can see that  $x_{0.9} = 90$  and also  $x_{0.95} = 90$ . Since the final 2% of probability is uniformly spread over the interval  $(90, 100)$ , we see that  $x_{0.99} = 95$ .

Again we condition on that part of the distribution which lies above the 90<sup>th</sup> percentile. The conditional distribution is now a mixed distribution, 0.08 of probability at  $x = 90$  and 0.02 spread uniformly over  $(90, 100)$ . Factoring these up, we have:

$$\begin{aligned}\Pr(X = 90) &= 0.8 \\ f(x) &= 0.02 \quad 90 < x \leq 100\end{aligned}$$

[Note that when the 0.02 of probability is originally spread over  $(90, 100)$ , it produces a PDF of 0.002. When this is factored up, it goes back to being 0.02.]

The mean of this distribution is:

$$0.8 \times 90 + 0.2 \times 95 = 91$$

Alternatively, we can use:

$$\text{TVaR}_\alpha(X) = \frac{(\beta - \alpha)x_\alpha + (1 - \beta)E[X | X > x_\alpha]}{1 - \alpha} = \frac{(0.98 - 0.9)90 + 0.02(95)}{0.1} = 91$$

using  $\alpha = 0.9$ ,  $\beta = 0.98$ , and  $E[X | X > x_\alpha] = 95$ .

Similarly, for  $\alpha = 0.95$ , we obtain a conditional distribution for which:

$$\begin{aligned}\Pr(X = 90) &= 0.6 \\ f(x) &= 0.04 \quad 90 < x \leq 100\end{aligned}$$

The mean of this conditional distribution is:

$$0.6 \times 90 + 0.4 \times 95 = 92$$

Again, the weighted average formula would also give us the correct answer.

If  $\alpha = 0.99$ , the 99<sup>th</sup> percentile does not lie at a discontinuity on the CDF. So we can just use  $E[X | X > 95]$ , which is the mean of a uniform distribution on the interval  $(95, 100)$ . The answer is 97.5.

**Solution 13.12**

The 99<sup>th</sup> percentile will be approximated by the 990<sup>th</sup> sample value, which will be 450.31.

[Alternatively, you could use 450.36, or a weighted average of these two figures].

The 99% CTE estimate is the mean of the figures in the last row of the table:

$$\bar{x} = \frac{1}{10}(450.36 + \dots + 767.20) = 556.52$$

---

**Solution 13.13**

The number of sample values below the 990<sup>th</sup> value will have a binomial distribution with parameters  $n = 1,000$  and  $\theta = 0.99$ . We will approximate this using a normal distribution with mean 990 and variance 9.9.

The distribution of the corresponding sample proportion will be normal with mean 0.990 and variance  $9.9 \times 10^{-6}$ .

So the confidence interval will be:

$$0.99 \pm 1.96\sqrt{9.9 \times 10^{-6}} = (0.98383, 0.99617)$$

So the interval in terms of sample values should run from the 983<sup>rd</sup> to the 997<sup>th</sup> value inclusive. This gives us a confidence interval of (405.47, 580.22).

---